



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY
WASHINGTON, DC 20460

OFFICE OF CHEMICAL SAFETY
AND POLLUTION PREVENTION

May 18, 2020

Beth Mitchell
Regulatory Affairs Manager
SePRO Corporation
11550 North Meridian Street, Suite 600
Carmel, IN 46032

Subject: PRIA Label Amendment – New Uses on Avocado, Tangerine (Mandarin),
Pomegranate, Pistachio, and Stone Fruit (Crop Group 12-12)
Product Name: BRAKE
EPA Registration Number: 67690-78
Application Date: 9/26/2018
Decision Number: 544749

Dear Ms. Mitchell:

The application referred to above, submitted under the Federal Insecticide, Fungicide, and Rodenticide Act (FIFRA), as amended is acceptable under FIFRA sec 3(c)(5). You must submit and/or cite all data required for registration/reregistration/registration review of your product when the Agency requires all registrants of similar products to submit such data.

A stamped copy of your labeling is enclosed for your records. This labeling supersedes all previously accepted labeling. You must submit one (1) copy of the final printed labeling before you release the product for shipment with the new labeling. In accordance with 40 CFR 152.130(c), you may distribute or sell this product under the previously approved labeling for 18 months from the date of this letter. After 18 months, you may only distribute or sell this product if it bears this new revised labeling or subsequently approved labeling. "To distribute or sell" is defined under FIFRA section 2(gg) and its implementing regulation at 40 CFR 152.3.

Should you wish to add/retain a reference to the company's website on your label, then please be aware that the website becomes labeling under FIFRA and is subject to review by the Agency. If the website is false or misleading, the product would be misbranded and unlawful to sell or distribute under FIFRA section 12(a)(1)(E). 40 CFR 156.10(a)(5) list examples of statements EPA may consider false or misleading. In addition, regardless of whether a website is referenced on your product's label, claims made on the website may not substantially differ from those claims approved through the registration process. Therefore, should the Agency find or if it is brought to our attention that a website contains false or misleading statements or claims substantially differing from the EPA approved registration, the website will be referred to the EPA's Office of Enforcement and Compliance.

Page 2 of 2
EPA Reg. No. 67690-78
Decision No. 544749

Your release for shipment of the product constitutes acceptance of these conditions. If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA section 6. If you have any questions, please contact Sarah Meadows by phone at 703-347-0505, or via email at meadows.sarah@epa.gov.

Sincerely,

A handwritten signature in black ink, appearing to read "D. Kenny", with a long horizontal flourish extending to the right.


Daniel Kenny, Chief
Herbicide Branch
Registration Division (7505P)
Office of Pesticide Programs

Enclosure

BRAKE®

EPA Reg. No. 67690-78

Sub-label A	Cotton	pp. 1-15
Sub-label B	Permanent Crops	pp. 16-28
	Extended Warranty Statement	pp. 29



[Sub-label A - Cotton: Front of label booklet]



FLURIDONE	GROUP	12	HERBICIDE
-----------	-------	----	-----------

Brake®

Preemergence herbicide for use in cotton to control pigweed.

Active Ingredient

Fluridone:

1-methyl-3-phenyl-5-[3-(trifluoromethyl)phenyl]-4(1*H*)-pyridinone 13.76%

Other Ingredients 86.24%

TOTAL 100.00%

Equivalent to 1.2 pounds fluridone per gallon.

KEEP OUT OF REACH OF CHILDREN

DANGER / PELIGRO

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail).

FIRST AID	
If in eyes	<ul style="list-style-type: none"> • Hold eye open and rinse slowly and gently with water for 15 to 20 minutes. • Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. • Call a poison control center or doctor for treatment advice.
If swallowed	<ul style="list-style-type: none"> • Call a poison control center or doctor immediately for treatment advice. • Have person sip a glass of water if able to swallow. • Do not induce vomiting unless told to do so by a poison control center or doctor. • Do not give anything by mouth to an unconscious person.
If inhaled	<ul style="list-style-type: none"> • Move person to fresh air. • If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth, if possible. • Call a poison control center or doctor for further treatment advice
HOTLINE NUMBER	
Have the product container or label with you when calling a poison control center or doctor, or going for treatment. In case of emergency endangering health or the environment involving this product, call INFOTRAC at 1-800-535-5053 .	

NOTE TO PHYSICIAN: Probable mucosal damage may contraindicate the use of gastric lavage.

SHAKE WELL BEFORE USING

Refer to label booklet for additional Precautionary Information and Directions for Use including Storage and Disposal.

NOTICE: Read the entire label before using. **Use only according to label directions. Before buying or using this product, read *Warranty Disclaimer* and *Misuse* statements in label booklet. If terms are unacceptable, return at once, unopened.**

Agricultural Chemical: Do not ship or store with food, feeds, drugs or clothing.

EPA Reg. No. 67690-78
FPL20200511

EPA Est. No. _____
[P/N]_____

Brake is a registered trademark of SePRO Corporation

SePRO Corporation • 11550 North Meridian Street, Suite 600 • Carmel, IN 46032, U.S.A.

Herbicide

Net contents _____ (Non-refillable)

[Sub-label A - Cotton: Label booklet]

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

Danger. Corrosive. Causes irreversible eye damage. Do not get in eyes or on clothing. Harmful if inhaled or swallowed. Avoid breathing spray mist. Prolonged or frequently repeated skin contact may cause allergic reactions in some individuals. Wear protective eyewear. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet. Remove and wash contaminated clothing before reuse.

PERSONAL PROTECTIVE EQUIPMENT (PPE)

Mixers, loaders, applicators and other handlers **must** wear:

- Protective eyewear;
- Long-sleeved shirt and long pants; and
- Shoes plus socks.

Discard clothing and other absorbent materials that have been drenched or heavily contaminated with this product's concentrate. Do not reuse them. Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

When handlers use closed systems or enclosed cabs in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40CFR 170.240(d)(4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

USER SAFETY RECOMMENDATIONS

Users should:

- Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

ENVIRONMENTAL HAZARDS

For terrestrial use only. **Toxic to fish and other aquatic organisms. Do not apply directly to water, or to areas where surface water is present, or to intertidal areas below the mean high water mark.** Do not contaminate water when disposing of equipment washwater or rinsate. Do not apply when weather conditions favor drift from target area.

Groundwater Advisory

This product has properties and characteristics associated with chemicals detected in groundwater. This chemical may leach into groundwater if used in areas where soils are permeable, particularly where the water table is shallow.

Surface Water Advisory

This product may impact surface water quality due to runoff of rain water. This is especially true for poorly draining soils and soils with shallow ground water. This product is classified as having high potential for reaching surface water via runoff for several months or more after application. A level, well-maintained vegetative buffer strip between areas to which this product is applied and surface water features such as ponds, streams, and springs will reduce the potential loading of fluridone from runoff water and sediment. While at least 0.5 inches of rainfall or irrigation is recommended after application, runoff of this product should be avoided and will be reduced by avoiding applications when runoff-producing rainfall is expected to occur within 48 hours.

Non-Target Organisms Advisory Statement

This product is toxic to plants and may adversely impact the forage and habitat of non-target organisms, including pollinators, in areas adjacent to the treated site. Protect the forage and habitat of non-target organisms by following label directions intended to minimize spray drift.

PHYSICAL AND CHEMICAL HAZARDS

Do not mix or allow contact with oxidizing agents. Hazardous chemical reaction may occur.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

DO NOT apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation. Ensure spray drift to non-target species does not occur.

Keep containers closed to avoid spills and contamination.

IN CASE OF EMERGENCY

In case of large-scale spillage regarding this product, call **INFOTRAC** at **1-800-535-5053**.

In case of medical emergency regarding this product, call:

- Your local doctor for immediate treatment;
- Your local poison control center (hospital);
- **INFOTRAC: 1-800-535-5053.**

Steps to be taken in case material is released or spilled:

- Dike and contain the spill with inert material (sand, earth, etc.) and transfer liquid and solid diking material to separate containers for disposal.
- Remove contaminated clothing, and wash affected skin areas with soap and water.
- Wash clothing before reuse.
- Keep the spill out of all sewers and open bodies of water.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE), restricted-entry interval, and notification to workers.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 48 Hours. Exception: If the product is soil-injected or soil-incorporated, the Worker Protection Standard, under certain circumstances, allows workers to enter the treated area if there will be no contact with anything that has been treated.

The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and involves contact with anything that has been treated, such as plants, soil, or water, is:

- Coveralls worn over short sleeved shirt and pants;
- Chemical resistant gloves;
- Shoes plus socks; and
- Protective eyewear.

PRODUCT INFORMATION

This product is a selective preemergence herbicide for use in cotton to control pigweed (e.g. *Palmer amaranth*). This product is also active on a range of annual grass and small seeded broadleaf weeds. Apply this product as preplant or preemergence surface application in coarse, medium or fine textured soils (refer to Table 2 for application rates for this product). Herbicidal symptoms of this product appear as white (chlorotic) or pink growing points in susceptible plant species. Cotton is tolerant to applications of this product when applied according to the label.

Resistance Management

This product contains a Group 12 herbicide. Any weed population may contain or develop plants naturally resistant to Group 12 herbicides. Combining active ingredients through tank mixing is one way to delay development of herbicide resistance. However, weed species with acquired resistance to Group 12 may eventually dominate the weed population if Group 12 herbicides are used repeatedly in the same field or in successive years as the primary method of control for targeted species. This may result in partial or total loss of control of those species by this product or other Group 12 herbicides.

Do not apply a product containing fluridone to the same field more than two years in a row. After the second year's application, use an alternative herbicide from Groups other than 12, effective in controlling the weeds of concern the following year.

To delay herbicide resistance, consider:

- Avoiding the consecutive use of this product and other target site of action Group 12 herbicides that have a similar target site of action, on the same weed species;
- Basing herbicide use on a comprehensive Integrated Pest Management (IPM) program;
- Monitoring treated weed populations for loss of field efficacy and treat weed escapes immediately with another group herbicide - start weed free and stay weed free;
- Contacting your local extension specialist, certified crop advisors, and/or manufacturer for herbicide resistance management and/or integrated weed management recommendations for specific crops and resistant weed biotypes;
- Consulting your agricultural dealer, consultant, applicator, and/or appropriate state agricultural extension service representative for specific alternative cultural practices or herbicide recommendations available in your area.
- Contacting a SePRO Corporation representative for further information or to report suspected resistance.

Integrated Pest Management

This product may be used as part of an Integrated Pest Management (IPM) program that can include biological, cultural, and genetic practices aimed at preventing economic pest damage. IPM principles and practices include field scouting or other detection methods, correct target pest identification, population monitoring, and treating when target pest populations reach locally determined action thresholds. Consult your state cooperative extension service, professional consultants or other qualified authorities to determine appropriate action treatment threshold levels for treating specific pest/crop systems in your area.

Use Restrictions

- **DO NOT** apply by air.
- Chemigation: **DO NOT** apply through any type of irrigation system.
- **DO NOT** apply more than 32 fluid ounces of this product per acre per year (equivalent to 0.30 lb fluridone per acre).
- **DO NOT** apply a product containing fluridone to the same field more than 2 years in a row.
- Apply prior to planting or within 36 hours after planting cotton.
- **DO NOT** apply this product over the top of cotton plants as cotton injury will likely occur.
- Organic fertilizer must be incorporated into soil prior to application of this product to the soil surface.

Use Precautions

- Thoroughly clean the spray system with water and a suitable tank cleaner before and after each use.
- To provide adequate coverage, it is recommended that ground speed not exceed 10 mph during application.
- Avoid overlapping spray swaths as injury may occur to rotational crops.

Rotational Crop Restrictions

This product may persist in the soil past the cotton growing season. To avoid difficulties in establishing rotational crops, follow the restrictions in Table 1:

Table 1. Rotational crop restrictions for Brake
--

Crop to be Planted	Minimum Rotation Interval (months after last Brake application) [†] ; ^{††}
Application rate of 16 fluid ounces Brake per acre	
Cotton	0
Soybean and sweet potato	2
Rice, peanut, wheat, barley, and rye	8 ^{†††}
Corn and sorghum	10 ^{††††}
Alfalfa, sunflower, and crops in the Solanaceae family (e.g. peppers, tobacco, tomatoes)	18
Application rates between 16.1 and 21 fluid ounces Brake per acre	
Cotton	0
Sweet potato	2
Soybean	4
Rice, peanut, wheat, barley, and rye	8
Corn and sorghum	12
Alfalfa, sunflower, and crops in the Solanaceae family (e.g. pepper, tobacco, tomatoes)	18
Application rates between 21.1 and 32 fluid ounces Brake per acre	
Cotton	0
Sweet potato	2
Peanut	8
Soybean, rice, wheat, barley, and rye	12
Alfalfa, corn, sorghum, sunflower, and crops in the Solanaceae family (e.g. peppers, tobacco, tomatoes)	18

[†] West TX only: Add 4 months to rotational restrictions for all crops except cotton and peanut.

^{††} Rotation intervals for Brake may be reduced if soil analyses indicate fluridone concentrations less than the level of detection (0.02 ug/g) using appropriate analytical techniques or through bioassay. Soil samples used for analysis must be representative of the field(s) in question and collected to a depth of 6 inches. User accepts all responsibility for any crop injury if rotation intervals are reduced.

^{†††} If used as a cover crop (i.e. will not be harvested), wheat, barley and rye may be planted 5 months after the last Brake application if no more than 16 fluid ounces per acre were applied. Minor chlorosis and stunting of the small grains may occur, especially if heavy rainfall occurs during or soon after emergence, but plants typically outgrow these effects and develop normally.

^{††††} Minor chlorosis and stunting of corn and sorghum may occur, especially if heavy rainfall occurs during or soon after emergence, but plants typically outgrow these effects and develop normally.

- If this product is used for two consecutive years in the same field at a rate of 16 fl oz per acre, only plant cotton, peanut, or soybean the following year in those fields.
- If this product is used for two consecutive years in the same field at a rate greater than 16 fl oz per acre, only plant cotton or peanut the following year in those fields.
- In the event of poor cotton stand due to adverse weather or other circumstances, a field treated with this product may be replanted with cotton following light tillage or planted in the undisturbed soil. During replanting, minimum tillage is recommended to preserve the herbicide barrier for effective weed control.

- If a field needs to be replanted with cotton due to poor stand or stand failure, do not apply a second application of this product. If tank-mix combinations were initially used, refer to product labels for any additional replanting instruction.
- Follow the crop rotation intervals specified in Table 1. For any crop not specifically listed in Table 1, do not replant for at least 18 months in fields treated with this product, as crop injury may occur.
- Over-application may result in crop injury or injury to rotational crops. Avoid overlapping spray swaths, as injury may occur to rotational crops.

MIXING AND APPLICATION DIRECTIONS

Apply this product evenly and uniformly to the soil surface with a minimum spray solution of 10 gallons per acre. Fill spray tank partially full with water (e.g. ½ full). Start agitation. Add correct quantity of this product. **Maintain continuous agitation while filling spray tank to required volume and during application. Do not stop agitation prior to or during application.** Nozzle selection should meet manufacturer's volume and pressure recommendations for preplant surface or preemergence applications. Uniform spray coverage is essential for good efficacy.

Tank Mix Directions

This product may be tank-mixed with other herbicides to enhance efficacy provided that the labels for this product and the tank-mix partner do not prohibit such mixing. **Due to the many herbicide formulations available and to ensure compatibility, ALWAYS conduct a jar test before field application of any tank-mix combination.** Do not allow tank-mixed material to sit in the spray tank for extended periods of time prior to application (e.g. overnight).

For all tank mixtures, it is the pesticide user's responsibility to ensure that all products are registered for the intended use. Read and follow the applicable restrictions and limitations and directions for use on all product labels involved in tank mixing. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture.

NOTE: Tank mixing or use of this product with any other product which is not specifically and expressly authorized by the label shall be at the exclusive risk of the user, applicator and/or application adviser, to the extent allowed by applicable law.

Spray Additives and Fertilizers: No adjuvants are needed for application of this product. However, if this product is part of a tank-mix that includes a burndown herbicide or combination, follow label use directions for the adjuvant/surfactant and other herbicides used in the tank-mix. Conduct a jar test before mixing this product with adjuvants and/or fertilizer to ensure compatibility. Generally, tank mix additives should be added last to the tank mix. Do not mix this product with spray adjuvants or drift control agents containing polyacrylamide.

Soil Preparation – Start Weed Free

This product will not control established weeds. It is important to **Start Weed Free** by controlling all existing weeds prior to and/or during this product's application. Depending on

the type of tillage, established weeds could be covered with soil and shielded from exposure to the burndown herbicide. Therefore, for best results, apply burndown herbicides 24 to 96 hours prior to cultivation in conventional and strip tillage systems. Depending on environmental conditions, tillage can also disturb the seed bank and cause more weeds to germinate. Therefore, it is recommended to tank-mix this product with a burndown herbicide, such as Gramoxone® (active ingredient paraquat labeled for use in cotton), at the time of planting. It is the pesticide user's responsibility to ensure that all products in the listed mixtures are registered for the intended use. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture.

APPLICATION RATES

Apply this product as a preplant surface or preemergence application. Refer to Table 2 for specific rates by soil type. Use lower rates in areas where weed pressure is low and higher rates in areas where weed pressure is heavy. Application of this product to soils with high clay content may require higher rates. Application of this product to soils with high organic matter (e.g. >3%) may result in reduced efficacy.

Table 2. Application rates for Brake [†]		
Soil Texture	Rate in Fluid Ounces per Acre	Application Method
All Soil Types	16 to 32	Preemergence; or preplant up to 14 days prior to planting

[†] If applying less than 21 fluid ounces per acre, another residual herbicide effective at controlling the target weed species must be tank mixed with this product. Use in accordance with the most restrictive label limitations and precautions of the products used in the tank-mix. Do not exceed any labeled rate. Due to the many herbicide formulations available and to ensure compatibility, always conduct a jar test before field application of any tank-mix combination.

Band Application

This product may be applied as a banded application. When this product is applied on a band, calculate the amount of herbicide and water volume needed for band treatment by the following formulas:

$$\frac{\text{Band width in inches}}{\text{Row width in inches}} \times \frac{\text{Broadcast rate}}{\text{per acre}} = \text{Band herbicide rate per acre}$$

$$\frac{\text{Band width in inches}}{\text{Row width in inches}} \times \frac{\text{Broadcast volume}}{\text{per acre}} = \text{Band water volume per acre}$$

Band width must not exceed 50% (one-half) the cotton row width. Best coverage is obtained with a minimum of two nozzles, one directed to each side of the planted row. Application with a single nozzle directed over the top of the row is suitable for preemergence applications. Cultivation of untreated areas may be needed following band applications.

APPLICATION METHODS

Germinating weeds can be controlled by soil residual activity from either preplant surface or preemergence applications of this product in no till, strip or conventional tillage situations.

Preemergence Applications

Apply this product as a preemergence treatment to coarse, medium or fine textured soils at rates from 16 to 32 fluid ounces per acre. **If applying less than 21 fluid ounces per acre, another residual herbicide effective at controlling the target weed species must be tank mixed with this product.** Use in accordance with the most restrictive label limitations and precautions of the products used in the tank-mix. Do not exceed any labeled rate and only tank mix with other herbicides labeled for use in cotton and compatible with the cotton trait being planted. For best results, apply behind the planter (i.e. at planting) or within 36 hours after planting. Do not apply more than 32 fluid ounces per acre. Moisture is necessary to activate this product in soil for residual weed control. Dry weather following applications of this product may reduce effectiveness. For improved weed control, a minimum of 0.5 inches of rain or overhead irrigation is recommended following application.

Preplant Surface Applications

This product may be applied as preplant surface application in coarse, medium or fine textured soils at rates from 16 to 32 fluid ounces per acre up to 14 days before planting. **If applying less than 21 fluid ounces per acre, another residual herbicide effective at controlling the target weed species must be tank mixed with this product.** Use in accordance with the most restrictive label limitations and precautions of the products used in the tank-mix. Do not exceed any labeled rate and only tank mix with other herbicides labeled for use in cotton and compatible with the cotton trait being planted. Do not apply more than 32 fluid ounces per acre. Apply after the last tillage operation is completed. A minimum of 0.5 inches of rain or overhead irrigation must occur before planting cotton, and the seedbed must not be disturbed or reworked following application.

Controlling Early Weed Escapes

DO NOT use this product as a stand-alone herbicide. This product must be used as part of a comprehensive weed control program. Weeds may escape control prior to this product being activated by moisture. **A postemergence application within 18 days after planting that includes glyphosate, glufosinate, or other postemergence herbicide that is effective at controlling the target weed species and compatible with the cotton variety, combined with a residual herbicide (e.g. Warrant®, Dual Magnum®, or Outlook®) IS REQUIRED to control early germinating weeds and to overlap residual herbicides.** The purpose of overlapping residual herbicides during postemergence applications is to eliminate all weed escapes for a zero-tolerance approach. Aggressive scouting is recommended and, when necessary, use secondary postemergence applications including a residual herbicide. More frequent postemergence applications may be necessary in locations with very sandy soils (>60% sand) due to the increased potential for rapid changes in soil moisture. Depending on the frequency of weed escapes, standard post-directed or layby herbicide applications are also recommended, when necessary, in addition to other cultural practices. **Only use herbicides that are compatible with the cotton variety being grown.** If uncertain, contact a local extension agent or crop consultant to verify herbicide tolerance for different cotton varieties.

The Brake Preemergence Four-Step System: *Start Weed Free. Stay Weed Free.*

1. Preplant burndown and/or cultivate;

2. Apply this product at planting or within 36 hours of planting with a burndown herbicide;
3. Apply a postemergence herbicide containing glyphosate, glufosinate, or other postemergence herbicide that is effective at controlling the target weed and compatible with the cotton variety, combined with a residual herbicide within 18 days after planting; and
4. Scout and utilize postemergence and layby applications with overlapping residuals as needed.

SPRAY DRIFT MANAGEMENT

Boom-less Ground Applications

Setting nozzles at the lowest effective height will help to reduce the potential for spray drift.

Handheld Technology Applications

Take precautions to minimize spray drift.

SPRAY DRIFT ADVISORIES

The applicator is responsible for avoiding off-site spray drift. Be aware of nearby non-target sites and environmental conditions.

Importance of Droplet Size

An effective way to reduce spray drift is to apply large droplets. Use the largest droplets that provide target pest control. While applying larger droplets will reduce spray drift, the potential for drift will be greater if applications are made improperly or under unfavorable environmental conditions.

Controlling Droplet Size – Ground Boom

- Volume - Increasing the spray volume so that larger droplets are produced will reduce spray drift. Use the highest practical spray volume for the application. If a greater spray volume is needed, consider using a nozzle with a higher flow rate.
- Pressure - Use the lowest spray pressure recommended for the nozzle to produce the target spray volume and droplet size.
- Spray Nozzle - Use a spray nozzle that is designed for the intended application. Consider using nozzles designed to reduce drift.

Boom Height – Ground Boom

For ground equipment, the boom should remain level with the crop and have minimal bounce.

Shielded Sprayers

Shielding the boom or individual nozzles can reduce spray drift. Consider using shielded sprayers. Verify that the shields are not interfering with the uniform deposition of the spray on the target area.

Temperature and Humidity

When making applications in hot and dry conditions, use larger droplets to reduce effects of evaporation.

Temperature Inversions

Drift potential is high during a temperature inversion. Temperature inversions are characterized by increasing temperature with altitude and are common on nights with limited cloud cover and light to no wind. The presence of an inversion can be indicated by ground fog or by the movement of smoke from a ground source or an aircraft smoke generator. Smoke that layers and moves laterally in a concentrated cloud (under low wind conditions) indicates an inversion, while smoke that moves upward and rapidly dissipates indicates good vertical air mixing. Avoid applications during temperature inversions.

Wind

Drift potential generally increases with wind speed. **AVOID APPLICATIONS DURING GUSTY WIND CONDITIONS.**

Applicators need to be familiar with local wind patterns and terrain that could affect spray drift.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.

Pesticide Storage: Keep from freezing. Store in original container only. Do not store near feed or foodstuffs. In case of leak or spill, use absorbent materials to contain liquids and dispose as waste.

Pesticide Disposal: Wastes resulting from use of this product may be used according to label directions or disposed of at an approved waste disposal facility.

Container Handling

Nonrefillable Container. DO NOT reuse or refill this container. Triple rinse or pressure rinse container (or equivalent) promptly after emptying; then offer for recycling, if available, or reconditioning, if appropriate, or puncture and dispose of in a sanitary landfill, or by incineration, or by other procedures approved by state and local authorities.

Triple rinse containers small enough to shake (capacity ≤ 5 gallons) as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container ¼ full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank, or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times.

Triple rinse containers too large to shake (capacity >5 gallons) as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container ¼ full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank, or store rinsate for later use or disposal. Repeat this procedure two more times.

Pressure rinse as follows: Empty the remaining contents into application equipment or mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix tank, or collect rinsate for later use or disposal. Insert pressure rinsing nozzle in the side of the container and rinse at about 40 PSI for at least 30 seconds. Drain for 10 seconds after the flow begins to drip.

Container Handling (bulk): Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by incineration, or by other procedures approved by state and local authorities.

Warranty Disclaimer: SePRO Corporation warrants that this product conforms to the chemical description on the product label. Testing and research have also determined that this product is reasonably fit for the uses described on the product label. To the extent consistent with applicable law, SePRO Corporation makes no other express or implied warranty of fitness or merchantability nor any other express or implied warranty and any such warranties are expressly disclaimed.

Misuse: Federal law prohibits the use of this product in a manner inconsistent with its label directions. To the extent consistent with applicable law, the buyer assumes responsibility for any adverse consequences if this product is not used according to its label directions. In no case shall SePRO Corporation be liable for any losses or damages resulting from the use, handling or application of this product in a manner inconsistent with its label.

For additional important labeling information regarding SePRO Corporation's Terms and Conditions of Use, Inherent Risks of Use and Limitation of Remedies, please visit <http://seprolabels.com/terms> or scan the image below.



© Copyright ____ SePRO Corporation

®Brake is a registered trademark of SePRO Corporation

® Warrant is a registered trademark of Monsanto.

®Dual and Gramoxone are registered trademarks of Syngenta Group Company.

®Outlook is a registered trademark of BASF Corporation.

[Sub-label A - Cotton: Base Label]



FLURIDONE	GROUP	12	HERBICIDE
-----------	-------	----	-----------

Brake®

Active Ingredient

Fluridone:

1-methyl-3-phenyl-5-[3-(trifluoromethyl)phenyl]-4(1*H*)-pyridinone 13.76%

Other Ingredients 86.24%

TOTAL 100.00%

Equivalent to 1.2 pounds fluridone per gallon.

KEEP OUT OF REACH OF CHILDREN

DANGER / PELIGRO

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail).

FIRST AID	
If in eyes	<ul style="list-style-type: none"> • Hold eye open and rinse slowly and gently with water for 15 to 20 minutes. • Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. • Call a poison control center or doctor for treatment advice.
If swallowed	<ul style="list-style-type: none"> • Call a poison control center or doctor immediately for treatment advice. • Have person sip a glass of water if able to swallow. • Do not induce vomiting unless told to do so by a poison control center or doctor. • Do not give anything by mouth to an unconscious person.
If inhaled	<ul style="list-style-type: none"> • Move person to fresh air. • If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth, if possible. • Call a poison control center or doctor for further treatment advice
HOTLINE NUMBER	
Have the product container or label with you when calling a poison control center or doctor, or going for treatment. In case of emergency endangering health or the environment involving this product, call INFOTRAC at 1-800-535-5053 .	

NOTE TO PHYSICIAN: Probable mucosal damage may contraindicate the use of gastric lavage.

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

Corrosive. Causes irreversible eye damage. Do not get in eyes or on clothing. Harmful if

inhaled or swallowed. Avoid breathing spray mist. Prolonged or frequently repeated skin contact may cause allergic reactions in some individuals. Wear protective eyewear. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet. Remove and wash contaminated clothing before reuse.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.

Pesticide Storage: Keep from freezing. Store in original container only. Do not store near feed or foodstuffs. In case of leak or spill, use absorbent materials to contain liquids and dispose as waste.

Pesticide Disposal: Wastes resulting from use of this product may be used according to label directions or disposed of at an approved waste disposal facility.

Container Handling

Nonrefillable Container. DO NOT reuse or refill this container. Triple rinse or pressure rinse container (or equivalent) promptly after emptying; then offer for recycling, if available, or reconditioning, if appropriate, or puncture and dispose of in a sanitary landfill, or by incineration, or by other procedures approved by state and local authorities.

See inside label for complete container disposal directions including triple rinsing and pressure rinsing instructions.

SHAKE WELL BEFORE USING

Refer to label booklet for additional Precautionary Information and Directions for Use.

NOTICE: Read the entire label before using. **Use only according to label directions. Before buying or using this product, read *Warranty Disclaimer* and *Misuse* statements in label booklet. If terms are unacceptable, return at once, unopened.**

Agricultural Chemical: Do not ship or store with food, feeds, drugs or clothing.

EPA Reg. No. 67690-78
FPL20200511

EPA Est. No. _____
[P/N]_____

Brake is a registered trademark of SePRO Corporation

SePRO Corporation • 11550 North Meridian Street, Suite 600 • Carmel, IN 46032, U.S.A.

Herbicide

Net contents _____ (Non-refillable)

[Sub-label B – Permanent Crops: Front of label booklet]



FLURIDONE	GROUP	12	HERBICIDE
-----------	-------	-----------	-----------

Brake®

Preemergence herbicide for the control of annual grass and broadleaf weeds in avocado, tangerine (mandarin), pomegranate, pistachio, and stone fruit.

Active Ingredient

Fluridone:

1-methyl-3-phenyl-5-[3-(trifluoromethyl)phenyl]-4(1*H*)-pyridinone 13.76%

Other Ingredients 86.24%

TOTAL 100.00%

Equivalent to 1.2 pounds fluridone per gallon.

KEEP OUT OF REACH OF CHILDREN

DANGER / PELIGRO

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail).

FIRST AID	
If in eyes	<ul style="list-style-type: none"> • Hold eye open and rinse slowly and gently with water for 15 to 20 minutes. • Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. • Call a poison control center or doctor for treatment advice.
If swallowed	<ul style="list-style-type: none"> • Call a poison control center or doctor immediately for treatment advice. • Have person sip a glass of water if able to swallow. • Do not induce vomiting unless told to do so by a poison control center or doctor. • Do not give anything by mouth to an unconscious person.
If inhaled	<ul style="list-style-type: none"> • Move person to fresh air. • If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth, if possible. • Call a poison control center or doctor for further treatment advice
HOTLINE NUMBER	
<p>Have the product container or label with you when calling a poison control center or doctor, or going for treatment. In case of emergency endangering health or the environment involving this product, call INFOTRAC at 1-800-535-5053.</p>	

NOTE TO PHYSICIAN: Probable mucosal damage may contraindicate the use of gastric lavage.

SHAKE WELL BEFORE USING

Refer to label booklet for additional Precautionary Information and Directions for Use including Storage and Disposal.

NOTICE: Read the entire label before using. **Use only according to label directions. Before buying or using this product, read *Warranty Disclaimer* and *Misuse* statements in label booklet. If terms are unacceptable, return at once, unopened.**

Agricultural Chemical: Do not ship or store with food, feeds, drugs or clothing.

EPA Reg. No. 67690-78
FPL20200511

EPA Est. No. _____
[P/N]_____

Brake is a registered trademark of SePRO Corporation

SePRO Corporation • 11550 North Meridian Street, Suite 600 • Carmel, IN 46032, U.S.A.

Herbicide

Net contents _____ (Non-refillable)

[Sub-label B – Permanent Crops: Label booklet]

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

Danger. Corrosive. Causes irreversible eye damage. Do not get in eyes or on clothing. Harmful if inhaled or swallowed. Avoid breathing spray mist. Prolonged or frequently repeated skin contact may cause allergic reactions in some individuals. Wear protective eyewear. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet. Remove and wash contaminated clothing before reuse.

PERSONAL PROTECTIVE EQUIPMENT (PPE)

Mixers, loaders, applicators and other handlers **must** wear:

- Protective eyewear;
- Long-sleeved shirt and long pants; and
- Shoes plus socks.

Discard clothing and other absorbent materials that have been drenched or heavily contaminated with this product's concentrate. Do not reuse them. Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

When handlers use closed systems or enclosed cabs in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40CFR 170.240(d)(4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

USER SAFETY RECOMMENDATIONS

Users should:

- Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

ENVIRONMENTAL HAZARDS

For terrestrial use only. **Toxic to fish and other aquatic organisms. Do not apply directly to water, or to areas where surface water is present, or to intertidal areas below the mean high water mark.** Do not contaminate water when disposing of equipment washwater or rinsate. Do not apply when weather conditions favor drift from target area.

Groundwater Advisory

This product has properties and characteristics associated with chemicals detected in groundwater. This chemical may leach into groundwater if used in areas where soils are permeable, particularly where the water table is shallow.

Surface Water Advisory

This product may impact surface water quality due to runoff of rain water. This is especially true for poorly draining soils and soils with shallow ground water. This product is classified as having high potential for reaching surface water via runoff for several months or more after application. A level, well-maintained vegetative buffer strip between areas to which this product is applied and surface water features such as ponds, streams, and springs will reduce the potential loading of fluridone from runoff water and sediment. While at least 0.5 inches of rainfall or irrigation is recommended after application, runoff of this product should be avoided and will be reduced by avoiding applications when runoff-producing rainfall is expected to occur within 48 hours.

Non-Target Organisms Advisory Statement

This product is toxic to plants and may adversely impact the forage and habitat of non-target organisms, including pollinators, in areas adjacent to the treated site. Protect the forage and habitat of non-target organisms by following label directions intended to minimize spray drift.

PHYSICAL AND CHEMICAL HAZARDS

Do not mix or allow contact with oxidizing agents. Hazardous chemical reaction may occur.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

DO NOT apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation. Ensure spray drift to non-target species does not occur.

Keep containers closed to avoid spills and contamination.

IN CASE OF EMERGENCY

In case of large-scale spillage regarding this product, call **INFOTRAC** at **1-800-535-5053**.

In case of medical emergency regarding this product, call:

- Your local doctor for immediate treatment;
- Your local poison control center (hospital);
- **INFOTRAC: 1-800-535-5053.**

Steps to be taken in case material is released or spilled:

- Dike and contain the spill with inert material (sand, earth, etc.) and transfer liquid and solid diking material to separate containers for disposal.
- Remove contaminated clothing, and wash affected skin areas with soap and water.
- Wash clothing before reuse.
- Keep the spill out of all sewers and open bodies of water.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides.

It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE), restricted-entry interval, and notification to workers.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 48 Hours.

The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and involves contact with anything that has been treated, such as plants, soil, or water, is:

- Coveralls worn over short sleeved shirt and pants;
- Chemical resistant gloves;
- Shoes plus socks; and
- Protective eyewear.

PRODUCT INFORMATION

This product is a preemergence herbicide for control of annual grass and broadleaf weeds in Avocado, Tangerine (Mandarin), Pomegranate, Pistachio, and Stone Fruit (Crop Group 12-12). This product controls weeds by inhibiting carotenoid biosynthesis. Herbicidal symptoms of this product appear as white (chlorotic) or pink growing points in susceptible weed species.

Resistance Management

This product contains a Group 12 herbicide. Any weed population may contain or develop plants naturally resistant to Group 12 herbicides. Combining active ingredients through tank mixing is one way to delay development of herbicide resistance. However, weed species with acquired resistance to Group 12 may eventually dominate the weed population if Group 12 herbicides are used repeatedly in the same field or in successive years as the primary method of control for targeted species. This may result in partial or total loss of control of those species by this product or other Group 12 herbicides.

Do not apply a product containing fluridone to the same orchard more than two years in a row. After the second year's application, use an alternative herbicide from Groups other than 12, effective in controlling the weeds of concern the following year.

To delay herbicide resistance, consider:

- Avoiding the consecutive use of this product and other target site of action Group 12 herbicides that have a similar target site of action, on the same weed species;
- Basing herbicide use on a comprehensive Integrated Pest Management (IPM) program;
- Monitoring treated weed populations for loss of field efficacy and treat weed escapes immediately with another group herbicide - start weed free and stay weed free;
- Contacting your local extension specialist, certified crop advisors, and/or manufacturer for herbicide resistance management and/or integrated weed management recommendations for specific crops and resistant weed biotypes;

- Consulting your agricultural dealer, consultant, applicator, and/or appropriate state agricultural extension service representative for specific alternative cultural practices or herbicide recommendations available in your area.
- Contacting a SePRO Corporation company representative for further information or to report suspected resistance.

Integrated Pest Management

This product may be used as part of an Integrated Pest Management (IPM) program that can include biological, cultural, and genetic practices aimed at preventing economic pest damage. IPM principles and practices include field scouting or other detection methods, correct target pest identification, population monitoring, and treating when target pest populations reach locally determined action thresholds. Consult your state cooperative extension service, professional consultants or other qualified authorities to determine appropriate action treatment threshold levels for treating specific pest/crop systems in your area.

Use Restrictions

- **DO NOT** apply by air.
- Chemigation: **DO NOT** apply through any type of irrigation system.
- **DO NOT** apply more than 43 fluid ounces of this product per acre per calendar year (equivalent to 0.40 lb fluridone per acre).
- Apply this product once per calendar year as a preemergence application to the soil surface.
- **DO NOT** apply a product containing fluridone to the same orchard or grove more than 2 years in a row.
- **DO NOT** apply if the soil is frozen or covered with snow.
- **DO NOT** apply within 30 days prior to harvest.
- **DO NOT** apply when fruit or nuts intended for harvest are on the ground as illegal residues may result.
- **DO NOT** apply using mechanically-pressurized handguns.

Use Precautions

- To minimize potential injury to newly transplanted trees, apply this product around trees that have been established for a minimum of one year after transplanting and are exhibiting normal growth and vigor.
- Crop injury may occur if this product is applied to trunks that are not fully callused mature brown bark or protected from spray contact by nonporous wraps, grow tubes, or waxed containers.
- Avoid spray contact with crop foliage, roots, or fruit as it may cause crop injury.
- Application to soil with open channels or depressions may allow this product to move into the root zone through open channels and cause crop injury.
- Thoroughly clean the spray system with water and a suitable tank cleaner before and after each use.
- To provide adequate coverage, it is recommended that ground speed not exceed 10 mph during application.
- For best results, incorporate organic fertilizer into soil prior to application of this product to the soil surface.

Rotational Crop Restrictions

This product is intended for use in perennial tree crops listed in this label. New orchards or groves of perennial tree crops may be established in a location previously treated with this product 18 months after application if the soil has been thoroughly mixed to a depth of at least 6 inches prior to planting, or prior to 18 months if a bioassay is conducted and does not result in any observed herbicide symptoms in the intended crop. Do not rotate to any annual crop other than cotton within 36 months after application of this product. If other herbicides have also been used, follow the most restrictive label for the crop rotation interval.

MIXING AND APPLICATION DIRECTIONS

Apply this product evenly and uniformly to the soil surface as a preemergence broadcast or band surface application in coarse, medium or fine textured soils, with a minimum spray volume of 10 gallons per acre (refer to Table 1 for application rates for this product). Fill spray tank partially full with water (e.g. ½ full). Start agitation. Add correct quantity of this product. **Maintain continuous agitation while filling spray tank to required volume and during application. Do not stop agitation prior to or during application.** Nozzle selection should meet manufacturer's volume and pressure recommendations for preemergence surface applications. Uniform spray coverage is essential for good efficacy.

Tank Mix Directions

This product may be tank-mixed with other herbicides to enhance efficacy provided that the labels for this product and the tank-mix partner do not prohibit such mixing. **Due to the many herbicide formulations available and to ensure compatibility, ALWAYS conduct a jar test before field application of any tank-mix combination.** Do not allow tank-mixed material to sit in the spray tank for extended periods of time prior to application (e.g. overnight).

For all tank mixtures, it is the pesticide user's responsibility to ensure that all products are registered for the intended use. Read and follow the applicable restrictions and limitations and directions for use on all product labels involved in tank mixing. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture.

NOTE: Tank mixing or use of this product with any other product which is not specifically and expressly authorized by the label shall be at the exclusive risk of the user, applicator and/or application adviser, to the extent allowed by applicable law.

Spray Additives and Fertilizers: No adjuvants are needed for application of this product. However, if this product is part of a tank-mix that includes a burndown herbicide or combination, follow label use directions for the adjuvant/surfactant and other herbicides used in the tank-mix. Conduct a jar test before mixing this product with adjuvants and/or fertilizer to ensure compatibility. Generally, tank mix additives should be added last to the tank mix. Do not mix this product with spray adjuvants or drift control agents containing polyacrylamide.

Soil Preparation – Start Weed Free

This product will not control established weeds. It is important to **Start Weed Free** by controlling all existing weeds prior to and/or during this product's application. Therefore, when weeds are emerged at application, it is recommended to tank-mix this product with a burndown herbicide. It is the pesticide user's responsibility to ensure that all products in the listed mixtures are registered for the intended use. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture.

APPLICATION RATES

Apply this product once per calendar year as a preemergence application to the soil surface. Refer to Table 1 for specific rates. Use lower rate in areas where weed pressure is low and higher rate in areas where weed pressure is heavy. Application of this product to soils with high clay content may require higher rate. Application of this product to soils with high organic matter (e.g. >3%) may result in reduced efficacy.

Soil Texture	Rate in Fluid Ounces per Acre	Application Method
All Soil Types	21 to 43	Preemergence

[†] If applying less than 32 fluid ounces per acre, another residual herbicide effective at controlling the target weed species must be tank mixed with this product. Use in accordance with the most restrictive label limitations and precautions of the products used in the tank-mix. Do not exceed any labeled rate.

Band Application

This product may be applied as a banded application. When this product is applied in a band, calculate the amount of herbicide and water volume needed for band treatment by the following formulas:

$$\frac{\text{Band width in inches}}{\text{Row width in inches}} \times \frac{\text{Broadcast rate}}{\text{per acre}} = \text{Band herbicide rate per acre}$$

$$\frac{\text{Band width in inches}}{\text{Row width in inches}} \times \frac{\text{Broadcast volume}}{\text{per acre}} = \text{Band water volume per acre}$$

APPLICATION METHODS

This product provides residual control of susceptible annual grass and broadleaf weed species when applied as a broadcast or banded treatment prior to weed seed germination. Only use in Avocado, Tangerine (Mandarin), Pomegranate, Pistachio, and Stone Fruit (Crop Group 12-12).

Stone Fruit Group 12-12 Including: Apricot; apricot, Japanese; capulin; cherry, black; cherry, Nanking; cherry, sweet; cherry, tart; Jujube, Chinese; nectarine; peach; plum; plum, American; plum, beach; plum, Canada; plum, cherry; plum, Chickasaw; plum, Damson; plum, Japanese; plum, Klamath; plum, prune; plumcot; sloe; cultivars, varieties, and/or hybrids of these.

Preemergence Applications

Moisture is necessary to activate this product in soil for residual weed control. Dry weather following application of this product may reduce effectiveness. A minimum of 0.5 inches of rain or overhead irrigation is recommended following application. For improved weed control, apply supplemental irrigation during the growing season to keep soil moist.

Apply this product as a preemergence treatment to the soil surface at rates from 21 to 43 fluid ounces per acre. Avoid spray contact with crop foliage, green bark, roots, or fruit as it may cause crop injury. **If applying less than 32 fluid ounces per acre, another residual herbicide effective at controlling the target weed species must be tank mixed with this product.** Use in accordance with the most restrictive label limitations and precautions of the products used in the tank-mix. Do not exceed any labeled rate and only tank mix with other herbicides labeled for the intended use.

SPRAY DRIFT MANAGEMENT

Boom-less Ground Applications

Setting nozzles at the lowest effective height will help to reduce the potential for spray drift.

Handheld Technology Applications

Take precautions to minimize spray drift.

SPRAY DRIFT ADVISORIES

The applicator is responsible for avoiding off-site spray drift. Be aware of nearby non-target sites and environmental conditions.

Importance of Droplet Size

An effective way to reduce spray drift is to apply large droplets. Use the largest droplets that provide target pest control. While applying larger droplets will reduce spray drift, the potential for drift will be greater if applications are made improperly or under unfavorable environmental conditions.

Controlling Droplet Size – Ground Boom

- Volume - Increasing the spray volume so that larger droplets are produced will reduce spray drift. Use the highest practical spray volume for the application. If a greater spray volume is needed, consider using a nozzle with a higher flow rate.
- Pressure - Use the lowest spray pressure recommended for the nozzle to produce the target spray volume and droplet size.
- Spray Nozzle - Use a spray nozzle that is designed for the intended application. Consider using nozzles designed to reduce drift.

Boom Height – Ground Boom

For ground equipment, the boom should remain level with the crop and have minimal bounce.

Shielded Sprayers

Shielding the boom or individual nozzles can reduce spray drift. Consider using shielded sprayers. Verify that the shields are not interfering with the uniform deposition of the spray on the target area.

Temperature and Humidity

When making applications in hot and dry conditions, use larger droplets to reduce effects of evaporation.

Temperature Inversions

Drift potential is high during a temperature inversion. Temperature inversions are characterized by increasing temperature with altitude and are common on nights with limited cloud cover and light to no wind. The presence of an inversion can be indicated by ground fog or by the movement of smoke from a ground source or an aircraft smoke generator. Smoke that layers and moves laterally in a concentrated cloud (under low wind conditions) indicates an inversion, while smoke that moves upward and rapidly dissipates indicates good vertical air mixing. Avoid applications during temperature inversions.

Wind

Drift potential generally increases with wind speed. **AVOID APPLICATIONS DURING GUSTY WIND CONDITIONS.**

Applicators need to be familiar with local wind patterns and terrain that could affect spray drift.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.

Pesticide Storage: Keep from freezing. Store in original container only. Do not store near feed or foodstuffs. In case of leak or spill, use absorbent materials to contain liquids and dispose as waste.

Pesticide Disposal: Wastes resulting from use of this product may be used according to label directions or disposed of at an approved waste disposal facility.

Container Handling

Nonrefillable Container. DO NOT reuse or refill this container. Triple rinse or pressure rinse container (or equivalent) promptly after emptying; then offer for recycling, if available, or reconditioning, if appropriate, or puncture and dispose of in a sanitary landfill, or by incineration, or by other procedures approved by state and local authorities.

Triple rinse containers small enough to shake (capacity ≤ 5 gallons) as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container ¼ full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank, or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times.

Triple rinse containers too large to shake (capacity >5 gallons) as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container ¼ full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back

and forth several times. Empty the rinsate into application equipment or a mix tank, or store rinsate for later use or disposal. Repeat this procedure two more times.

Pressure rinse as follows: Empty the remaining contents into application equipment or mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix tank, or collect rinsate for later use or disposal. Insert pressure rinsing nozzle in the side of the container and rinse at about 40 PSI for at least 30 seconds. Drain for 10 seconds after the flow begins to drip.

Container Handling (bulk): Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by incineration, or by other procedures approved by state and local authorities.

Warranty Disclaimer: SePRO Corporation warrants that this product conforms to the chemical description on the product label. Testing and research have also determined that this product is reasonably fit for the uses described on the product label. To the extent consistent with applicable law, SePRO Corporation makes no other express or implied warranty of fitness or merchantability nor any other express or implied warranty and any such warranties are expressly disclaimed.

Misuse: Federal law prohibits the use of this product in a manner inconsistent with its label directions. To the extent consistent with applicable law, the buyer assumes responsibility for any adverse consequences if this product is not used according to its label directions. In no case shall SePRO Corporation be liable for any losses or damages resulting from the use, handling or application of this product in a manner inconsistent with its label.

For additional important labeling information regarding SePRO Corporation's Terms and Conditions of Use, Inherent Risks of Use and Limitation of Remedies, please visit <http://seprolabels.com/terms> or scan the image below.



© Copyright ____ SePRO Corporation

®Brake is a registered trademark of SePRO Corporation

[Sub-label B – Permanent Crops: Base Label]



FLURIDONE	GROUP	12	HERBICIDE
-----------	-------	----	-----------

Brake®

Active Ingredient

Fluridone:

1-methyl-3-phenyl-5-[3-(trifluoromethyl)phenyl]-4(1*H*)-pyridinone 13.76%

Other Ingredients 86.24%

TOTAL 100.00%

Equivalent to 1.2 pounds fluridone per gallon.

KEEP OUT OF REACH OF CHILDREN

DANGER / PELIGRO

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail).

FIRST AID	
If in eyes	<ul style="list-style-type: none"> • Hold eye open and rinse slowly and gently with water for 15 to 20 minutes. • Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. • Call a poison control center or doctor for treatment advice.
If swallowed	<ul style="list-style-type: none"> • Call a poison control center or doctor immediately for treatment advice. • Have person sip a glass of water if able to swallow. • Do not induce vomiting unless told to do so by a poison control center or doctor. • Do not give anything by mouth to an unconscious person.
If inhaled	<ul style="list-style-type: none"> • Move person to fresh air. • If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth, if possible. • Call a poison control center or doctor for further treatment advice
HOTLINE NUMBER	
Have the product container or label with you when calling a poison control center or doctor, or going for treatment. In case of emergency endangering health or the environment involving this product, call INFOTRAC at 1-800-535-5053 .	

NOTE TO PHYSICIAN: Probable mucosal damage may contraindicate the use of gastric lavage.

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

Corrosive. Causes irreversible eye damage. Do not get in eyes or on clothing. Harmful if

inhaled or swallowed. Avoid breathing spray mist. Prolonged or frequently repeated skin contact may cause allergic reactions in some individuals. Wear protective eyewear. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet. Remove and wash contaminated clothing before reuse.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.

Pesticide Storage: Keep from freezing. Store in original container only. Do not store near feed or foodstuffs. In case of leak or spill, use absorbent materials to contain liquids and dispose as waste.

Pesticide Disposal: Wastes resulting from use of this product may be used according to label directions or disposed of at an approved waste disposal facility.

Container Handling

Nonrefillable Container. DO NOT reuse or refill this container. Triple rinse or pressure rinse container (or equivalent) promptly after emptying; then offer for recycling, if available, or reconditioning, if appropriate, or puncture and dispose of in a sanitary landfill, or by incineration, or by other procedures approved by state and local authorities.

See inside label for complete container disposal directions including triple rinsing and pressure rinsing instructions.

SHAKE WELL BEFORE USING

Refer to label booklet for additional Precautionary Information and Directions for Use.

NOTICE: Read the entire label before using. **Use only according to label directions. Before buying or using this product, read *Warranty Disclaimer* and *Misuse* statements in label booklet. If terms are unacceptable, return at once, unopened.**

Agricultural Chemical: Do not ship or store with food, feeds, drugs or clothing.

EPA Reg. No. 67690-78
FPL20200511

EPA Est. No. _____
[P/N]_____

Brake is a registered trademark of SePRO Corporation

SePRO Corporation • 11550 North Meridian Street, Suite 600 • Carmel, IN 46032, U.S.A.

Herbicide

Net contents _____ (Non-refillable)

[Text accessed through the weblink / QR code. This is NOT part of the printed label]

TERMS AND CONDITIONS OF USE

If terms of the *Warranty Disclaimer* and *Misuse* provisions on the product label as well as the *Inherent Risks of Use* and *Limitation of Remedies* statements below are not acceptable, return unopened package at once to the seller for a full refund of purchase price paid. Otherwise, to the extent consistent with applicable law, use by the buyer or any other user constitutes acceptance of the terms under *Warranty Disclaimer*, *Misuse*, *Inherent Risks of Use*, and *Limitation of Remedies*.

INHERENT RISKS OF USE

It is impossible to eliminate all risks associated with use of this product. Plant injury, lack of performance, or other unintended consequences may result because of such factors as use of the product contrary to label instructions (including use under conditions noted on the label such as unfavorable temperatures, soil conditions, etc.), abnormal conditions (such as excessive rainfall, drought, tornadoes, hurricanes), the presence of other materials, the manner of application, or other factors, all of which are beyond the control of SePRO Corporation or the seller. To the extent consistent with applicable law, all such risks shall be assumed by the buyer and/or user of the product.

LIMITATION OF REMEDIES

To the extent consistent with applicable law, the exclusive remedy for losses or damages resulting from this product (including claims based on contract, negligence, strict liability, or other legal theories) shall be limited to, at SePRO Corporation's election, one of the following:

1. Refund of purchase price paid by buyer or user for product bought, or
2. Replacement of amount of product used.

To the extent consistent with applicable law, SePRO Corporation shall not be liable for losses or damages resulting from handling or use of this product unless SePRO Corporation is promptly notified of such losses or damages in writing. In no case shall SePRO Corporation be liable for consequential or incidental damages or losses.

The terms of the *Warranty Disclaimer* and *Misuse* provisions on the product label and these *Terms and Conditions of Use*, *Inherent Risks of Use* and *Limitation of Remedies* cannot be varied by any written or verbal statements or agreements. No employee or sales agent of SePRO Corporation or the seller is authorized to vary or exceed the terms of the *Warranty Disclaimer* and *Misuse* provisions on the product label and these *Terms and Conditions of Use*, *Inherent Risks of Use* and *Limitation of Remedies* in any manner.